arXiv:1207.4983v1 [math.PR] 20 Jul 2012

MINIMAL SPECTRAL REPRESENTATIONS OF INFINITELY DIVISIBLE AND MAX–INFINITELY DIVISIBLE PROCESSES

ZAKHAR KABLUCHKO AND STILIAN STOEV

Abstract. We introduce the notion of minimality for spectral representations of sum– and max–infinitely divisible processes and prove that the minimal spectral representation on a Borel space exists and is unique. This fact is used to show that a stationary, stochastically continuous, sum– or max–i.d. random process on $\mathbb{R}^d$ can be generated by a measure–preserving flow on a σ–finite Borel measure space and that this flow is unique. As a particular case, we characterize stationary, stochastically continuous, union–infinitely divisible random subsets of $\mathbb{R}^d$. We introduce several new classes of max–i.d. random fields including fields of Penrose type and fields associated to Poisson line processes.

1. Introduction and statement of results

1.1. **Introduction.** A stochastic process $X = \{X(t), t \in T\}$ is called *infinitely divisible* (i.d.) if for every $n \in \mathbb{N}$ it can be represented as a sum of n independent identically distributed (i.i.d.) processes. For finite–dimensional i.d. random vectors such classical concepts as Lévy measure and the Lévy–Itô decomposition are well known and play an important role; see [31]. It is natural to expect that these notions should have an analogue in the infinite–dimensional setting of i.d. stochastic processes. Pioneering work in this direction is due to Maruyama [20]. For any infinitely divisible process X he constructed a Lévy measure as a measure on the space $\mathbb{R}^T$ endowed with a certain σ–algebra. If the set T is countable, the Lévy measure enjoys a number of desirable properties, for example, it is σ–finite. In the particular case of stationary i.d. processes on $\mathbb{Z}$, Roy [28] used Maruyama's construction to establish a fruitful link with the infinite ergodic theory [1] and to generalize results known previously for stable processes [26, 27, 30], a subclass of the i.d. processes.

However, if the set T is not countable, the Lévy measure is not well–behaved. For example, it may be not σ–finite. The single point sets in the space $\mathbb{R}^T$ are not measurable with respect to the product σ–algebra on $\mathbb{R}^T$, so that even making a statement like "the Lévy measure does not charge the origin" becomes a non-trivial task. In the setting of stationary i.d. processes on $\mathbb{R}^d$ the natural shift transformations acting on the space $\mathbb{R}^{\mathbb{R}^d}$ leave the Lévy measure invariant, but they do not form a measurable group of transformations. All these facts prevent any straightforward generalization of the results of Roy [28] to the continuous–time setting, since ergodic theory usually requires the σ–finiteness of the invariant measure and the measurability of the flow. Maruyama [20]

2010 *Mathematics Subject Classification.* Primary, 60E07, 60H05; Secondary, 60G51, 60G52, 60G70, 60G60, 28D05.

Key words and phrases. Infinitely divisible process, max–infinitely divisible process, spectral representation, minimality, measure–preserving flow, Poisson process, stochastic integral.

Stilian A. Stoev was partially supported by the NSF grant DMS–1106695 at the University of Michigan.

gave a modification of his construction which makes the Lévy measure σ–finite, see Proposition 3.1 in [20], but it seems that there was no further investigation of this question since then.

In the present work we will give a construction of spectral representations of i.d. processes in the setting of uncountable index set T. This construction is different from Maruyama's one, but it can still be thought of as an adequate analogue of the Lévy measure and the Lévy–Itô decomposition. Our construction includes as a special case the spectral representations of stable processes [11, 26, 27]. We prove that the spectral representation of an i.d. process is *unique* provided one imposes a certain natural *minimality* condition. This result is then used to show that a stationary i.d. process can be generated as a stochastic integral over a measure–preserving flow. This generalizes some results of Rosiński [26, 27] on the spectral representations of stationary stable processes. The i.d. theory we develop here is even simpler (although it is more general) than the stable theory of [26, 27] since the i.d. theory involves dealing with *measure–preserving* flows instead of *non–singular* flows. The i.d. theory is not obtained as a straightforward generalization of the stable one. As was recently observed by Roy [29], in order to pass from a non–singular flow generating a stationary stable process to the measure–preserving flow generating the same process regarded as an i.d. process, one has to apply the construction of Maharam [19] from ergodic theory. In the stable case, the uniqueness of the minimal spectral representation is usually proved using the rigidity of L^α–spaces [10, 11, 26]. In the setting of i.d. processes, these methods are not available. Instead, we prove a general result on the existence of conjugacy between equimeasurable families of functions; see Lemma 3.1, below.

Our results have natural counterparts in the setting of *max–infinitely divisible* (max–i.d.) processes. A random vector is called max–i.d. if it can be represented as a maximum (taken componentwise) of n i.i.d. random vectors for every $n \in \mathbb{N}$. A similar definition applies to stochastic processes. Every one–dimensional distribution is max–i.d., but the situation changes in higher dimensions. Two–dimensional max–i.d. distributions were introduced by Balkema and Resnick [3], the general d–dimensional case was considered by Gerritse [7] and Vatan [34] (the latter work studies also max–i.d. vectors with values in $\mathbb{R}^\infty$). Representations in terms of suprema over a Poisson random measure were obtained by Giné et al. [8] for max–i.d. processes with continuous sample paths and by Balkema et al. [2] for stochastically continuous processes. In [33, 15, 35, 6] criteria for ergodicity and mixing of stationary max–i.d. processes have been given. For important results on max–stable processes (which form a subclass of the max–i.d. processes), see [4, 5, 33]. In the present work we will introduce the notion of minimality for spectral representations of max–i.d. processes and show that under this condition the appropriately modified spectral representation of Balkema et al. [2] is unique.

The paper is organized as follows. In Section 1.2 we introduce minimal spectral representations for max–i.d. processes and show their existence and uniqueness under the general Condition S of separability in probability. Section 1.3 contains parallel results for i.d. processes. The developed theory of minimal spectral representations is used to associate stationary i.d. and max–i.d. processes to measure preserving flows. This leads to extensions of existing decomposition and classification results for stable and max-stable processes. In Section 2 we present several examples and applications. For simplicity, we focus on the max–i.d. case. Analogous examples and results are valid for i.d. processes. The connection between the newly introduced minimal spectral representations and the existing ones is demonstrated Section 2.1. In Sections 2.2–2.5 we present new examples of stationary max–i.d. processes associated with dissipative and null flows. In Section 2.6 we characterize the stationary union infinitely divisible random sets by relating them to measure–preserving flows. The proofs are given in Section 3.

1.2. **Spectral representations of max–i.d. processes.** A stochastic process $X = \{X(t), t \in T\}$ defined on an index set T and taking values in $\mathbb{R}$ is called max–i.d., if for all $n \in \mathbb{N}$, there exist independent identically distributed (i.i.d.) processes $\{X_{i,n}(t),\ t \in T\}$, $i = 1, \dots, n$, such that

$$\{X(t),\ t \in T\} \overset{d}{=} \left\{ \max_{1 \le i \le n} X_{i,n}(t),\ t \in T \right\}. \tag{1}$$

Here, $\overset{d}{=}$ denotes the equality of finite–dimensional distributions. If $\{X(t), t \in T\}$ is a max–i.d. process, then for every collection of non–decreasing functions $\varphi_t : \mathbb{R} \to \mathbb{R}$, $t \in T$, the process $\{\varphi_t(X(t)), t \in T\}$ is also max–i.d. By choosing

$$\varphi_t(x) = \begin{cases} e^x, & \text{if } \operatorname{essinf} X(t) = -\infty, \\ x - \operatorname{essinf} X(t), & \text{if } \operatorname{essinf} X(t) > -\infty \end{cases}$$

we can always achieve that $\operatorname{essinf} \varphi_t(X(t)) = 0$. In the sequel we therefore assume without loss of generality that $\operatorname{essinf} X(t) = 0$ for every $t \in T$.

Balkema et al. [2] gave a representation of max–i.d. processes in terms of stochastic max–integrals over a Poisson random measure. This representation is, in general, non–unique; see Example 1.9 below. We will introduce the notion of *minimality* for representations of max–i.d. processes and prove that the minimal representation exists and is unique.

We recall the construction of Balkema et al. [2] in a form which is suitable for our purposes. Let $(\Omega, \mathcal{B}, \mu)$ be a σ–finite measure space. We denote by $\mathcal{L}^\vee = \mathcal{L}^\vee(\Omega, \mathcal{B}, \mu)$ the space of all measurable functions $f : \Omega \to \mathbb{R}$ such that $f \ge 0$ μ–a.e. and $\mu\{\omega : f(\omega) > a\}$ is finite for all $a > 0$. As usual, two functions are identified if they differ on a set of measure zero. Note that for every $f_1, f_2 \in \mathcal{L}^\vee$ and $c_1, c_2 \ge 0$ we have $\max(c_1 f_1, c_2 f_2) \in \mathcal{L}^\vee$. Next let us recall the definition of the max–integral from [2]. Let $\Pi_\mu = \{U_i, i \in J\}$ be a Poisson random measure on the space $(\Omega, \mathcal{B})$ with intensity μ. Here, J is at most countable index set. For $f \in \mathcal{L}^\vee$ define the stochastic max–integral

$$I(f) \equiv \int_\Omega^\vee f d\Pi_\mu := \sup_{i \in J} f(U_i). \tag{2}$$

Here, the supremum is taken over all atoms U_i of the Poisson process Π_μ. If Π_μ is empty, which can happen if $\mu(\Omega) < \infty$, then the supremum in the right–hand side is defined to be 0. From (2) one readily derives a formula for the joint distribution of the stochastic max–integrals: For all $f_1, \dots, f_n \in \mathcal{L}^\vee$ and $x_1, \dots, x_n \ge 0$ (not all of which are 0) we have

$$\begin{aligned} \mathbb{P}\{I(f_j) < x_j,\ 1 \le j \le n\} &= \mathbb{P}\Big\{\Pi_\mu\Big(\cup_{j=1}^n \{f_j \ge x_j\}\Big) = 0\Big\} \\ &= \exp\Big\{-\mu\Big(\cup_{j=1}^n \{f_j \ge x_j\}\Big)\Big\}. \end{aligned} \tag{3}$$

Observe that for any collection of deterministic functions $f_t \in \mathcal{L}^\vee,\ t \in T$, the process $\{I(f_t), t \in T\}$, is max–i.d. since the $X_{i,n}$'s in (1) can be defined by using independent copies of the same stochastic max–integrals but with respect to a Poisson point process with intensity $\frac{1}{n}\mu$.

Definition 1.1. Let $X = \{X(t), t \in T\}$ be a max–i.d. process with $\operatorname{essinf} X(t) = 0$ for all $t \in T$. A collection of functions $\{f_t, t \in T\} \subset \mathcal{L}^\vee(\Omega, \mathcal{B}, \mu)$ is a *spectral representation* of the process X if we have the following equality of laws:

$$\{X(t), t \in T\} \overset{d}{=} \left\{ \int_\Omega^\vee f_t d\Pi_\mu, t \in T \right\}, \tag{4}$$

where Π_μ is a Poisson random measure on $(\Omega, \mathcal{B})$ with intensity μ.

Here, we focus on the general class of processes that are separable in probability in the sense of the following definition.

Definition 1.2. A stochastic process $\{X(t), t \in T\}$ satisfies *Condition S* if there is at most countable set $T_0 \subset T$ such that for all $t \in T$, there exists a sequence $\{t_n\}_{n \in \mathbb{N}} \subset T_0$, with $X(t_n) \to X(t)$ in probability.

Theorem 1.3 (Balkema et al. [2])**.** *Let $\{X(t), t \in \mathbb{R}^d\}$ be a max–i.d. process satisfying Condition S. There exists a spectral representation of X defined on $\mathbb{R}$ endowed with the Lebesgue measure.*

We will prove existence and uniqueness of the spectral representation under the following condition of minimality.

Definition 1.4. A spectral representation $\{f_t, t \in T\} \subset \mathcal{L}^\vee(\Omega, \mathcal{B}, \mu)$ is called *minimal* if the following two conditions hold:

1. The σ–algebra generated by $\{f_t, t \in T\}$ coincides with $\mathcal{B}$ up to μ–zero sets. That is, for every $B \in \mathcal{B}$, exists an $A \in \sigma\{f_t, t \in T\}$, such that $\mu(A \Delta B) = 0$.
2. There is no set $B \in \mathcal{B}$ such that $\mu(B) > 0$ and for every $t \in T$, $f_t = 0$ a.e. on B.

Remark 1.5. The first condition does not imply the second one: consider $\Omega = \{0, 1\}$ with counting measure, $T = \{1\}$, and $f_1(\omega) = \omega$.

Theorem 1.6. *Let $X = \{X(t), t \in T\}$ be a max–i.d. process satisfying Condition S. There exists a minimal spectral representation of X defined on $[0, 1]$ endowed with a σ–finite Borel measure.*

We recall next several notions of isomorphisms from measure theory. For more details, see e.g. Ch. 22 in [32] and p. 167 in [9].

Definition 1.7. *(i)* An *isomorphism* between two *measurable spaces* $(\Omega_i, \mathcal{B}_i)$, $i = 1, 2$, is a bijection $\Phi : \Omega_1 \to \Omega_2$ such that both Φ and Φ^{-1} are measurable.

(ii) A measurable space $(\Omega, \mathcal{B})$ is said to be a *Borel space* if it is isomorphic (in the sense of part (i)) to a complete separable metric space endowed with the Borel σ–algebra. It is known that any Borel space is isomorphic to the interval $[0, 1]$ endowed with the Borel σ–algebra or to an at most countable set endowed with the σ–algebra of all subsets.

(iii) A Borel space endowed with a σ–finite measure will be called a *σ–finite Borel space.*

(iv) An *isomorphism (modulo null sets)* between two *measure spaces* $(\Omega_i, \mathcal{B}_i, \mu_i)$, $i = 1, 2$, is a bijection $\Phi : \Omega_1 \backslash A_1 \to \Omega_2 \backslash A_2$, where $A_1 \in \mathcal{B}_1$ and $A_1 \in \mathcal{B}_2$ are null sets, such that both Φ and Φ^{-1} are measurable and $\mu_1(A) = \mu_2(\Phi(A))$, for all measurable $A \subset \Omega_1 \backslash A_1$. Two isomorphisms Φ, Ψ are considered as equal modulo null sets if $\Phi(\omega) = \Psi(\omega)$ for μ_1–a.a. $\omega \in \Omega_1$.

The next theorem is the main result in this section. It establishes the uniqueness of the minimal spectral representation.

Theorem 1.8. *Let* $X = \{X(t), t \in T\}$ *be a max–i.d. process. If* $\{f_t^{(i)}, t \in T\}$ *are two minimal spectral representations of* X *defined on* σ*–finite Borel spaces* $(\Omega_i, \mathcal{B}_i, \mu_i)$*,* $i = 1, 2$*, then there is a measure space isomorphism* $\Phi : \Omega_1 \to \Omega_2$ *(defined modulo null sets) such that for all* $t \in T$*,*

$$f_t^{(1)}(\omega) = f_t^{(2)} \circ \Phi(\omega) \quad \textit{for } \mu_1 - a.a.\ \omega \in \Omega_1. \tag{5}$$

The measure space isomorphism Φ *with this property is unique modulo null sets.*

Example 1.9. Our definition of the space $\mathcal{L}^\vee$ of integrands is more restrictive than that of Balkema et al. [2], who allow measurable functions $f : \Omega \to \mathbb{R}$ with $\mu\{f \geq a\} < \infty$, for some $a \in \mathbb{R}$ (and don't assume that $\operatorname{essinf} X_t = 0$). With the definition used in [2] the uniqueness may fail, even for minimal representations. Let $\Omega_1 = \Omega_2 = \mathbb{Z}$ be endowed with the counting measure, let $T = \mathbb{Z} \cup \{*\}$ and define

$$f_t^{(1)}(\omega) = f_t^{(2)}(\omega) = \mathbb{1}_{\{t\}}(\omega) \text{ if } t \neq *, \quad f_*^{(1)}(\omega) = 1, \quad f_*^{(2)}(\omega) = \mathbb{1}_{\{\omega > 0\}} + \frac{1}{2}\mathbb{1}_{\{\omega \leq 0\}}.$$

One verifies readily that $\{f_t^{(i)}, t \in T\}$, $i = 1, 2$, are minimal representations of the same max–i.d. process. However, there is no bijection $\Phi : \Omega_1 \to \Omega_2$ such that $f_*^{(1)} \circ \Phi = f_*^{(2)}$. Note that $f_*^{(2)} \notin \mathcal{L}^\vee$ and hence Theorem 1.8 does not apply.

1.3. **Spectral representations of i.d. processes.**

A process $\{X(t),\ t \in T\}$ is said to be infinitely divisible (i.d.) if for all $n \in \mathbb{N}$ it can be represented (in distribution) as a sum of n independent and identically distributed processes. There is already extensive literature on the existence of spectral representations and the properties of i.d. processes (see e.g. [20, 28, 25] among many others). However, the minimality and the uniqueness of the spectral representations has not been considered so far.

Let $(\Omega, \mathcal{B}, \mu)$ be a $\sigma-$finite measure space. The space of integrands $\mathcal{L}^+$ consists of all measurable $f : \Omega \to \mathbb{R}$ such that

$$\int_\Omega \min\{\varepsilon, |f(\omega)|^2\}\mu(d\omega) < +\infty, \tag{6}$$

for some (or, equivalently, any) $\varepsilon > 0$. Functions differing on a set of measure 0 are identified. Observe that $\mathcal{L}^+$ is a linear space since $1 \wedge (f+g)^2 \leq 2(1 \wedge f^2 + 1 \wedge g^2)$. Following Maruyama [20], for $f \in \mathcal{L}^+$ define the stochastic integral

$$I(f) \equiv \int_\Omega^+ f d\Pi_\mu := \lim_{\varepsilon \to 0+} \Big\{ \sum_{i \in J} f(U_i)\mathbb{1}_{\{|f(U_i)| > \varepsilon\}} - \int_{\{|f| > \varepsilon\}} a(f) d\mu \Big\}, \tag{7}$$

where $\Pi_\mu = \{U_i, i \in J\}$ is a Poisson random measure on $(\Omega, \mathcal{B})$ with intensity μ and

$$a(u) = \begin{cases} u, & |u| \leq 1, \\ 1, & u > 1, \\ -1, & u < -1. \end{cases} \tag{8}$$

Note that the limit in (7) exists in the a.s. sense by the convergence theorem for L^2–bounded martingales. For $f_1, \ldots, f_n \in \mathcal{L}^+$ the joint distribution of the $I(f_j)$'s is characterized as follows. For all $\theta_1, \ldots, \theta_n \in \mathbb{R}$ we have

$$\mathbb{E}e^{i\sum_{j=1}^n \theta_j I(f_j)} = \exp\Big\{ \int_\Omega \Big(e^{i\sum_{j=1}^n \theta_j f_j(\omega)} - 1 - i\sum_{j=1}^n \theta_j a(f_j(\omega)) \Big) \mu(d\omega) \Big\}. \tag{9}$$

In particular, it is easy to verify that for all $f, g \in \mathcal{L}^+$ and $c \in \mathbb{R}$ we have $I(f+g) = I(f)+I(g)+\mathrm{const}$ and $I(cf) = cI(f) + \mathrm{const}$, i.e. the functional I is essentially linear up to additive constants (see also (30) below).

Definition 1.10. Let $X = \{X(t), t \in T\}$ be an i.d. process with trivial Gaussian component defined on some index set T. A collection of functions $\{f_t, t \in T\} \subset \mathcal{L}^+(\Omega, \mathcal{B}, \mu)$ is a spectral representation of the process X if we have the following equality of laws:

$$\{X(t), t \in T\} \stackrel{d}{=} \left\{ \int_\Omega^+ f_t d\Pi_\mu + c(t), t \in T \right\}, \tag{10}$$

where Π_μ is a Poisson random measure on $(\Omega, \mathcal{B}, \mu)$ and $c : T \to \mathbb{R}$ is some function.

Theorem 1.11. *Let $\{X(t), t \in T\}$ be an i.d. process which has a trivial Gaussian component and satisfies Condition S. There exists a minimal spectral representation of X defined on $[0, 1]$ endowed with a σ–finite Borel measure.*

Here, a spectral representation of an i.d. process is called minimal if it satisfies both conditions of Definition 1.4. To prove the theorem we will introduce the following truncated L^2–metric on the space $\mathcal{L}^+$:

$$d(f, g) \equiv d(f - g) := \Big(\int_\Omega 1 \wedge (f - g)^2 d\mu \Big)^{1/2}, \quad f, g \in \mathcal{L}^+. \tag{11}$$

Note that the triangle inequality follows from $1 \wedge |f - g| \le 1 \wedge |f| + 1 \wedge |g|$ and the triangle inequality in $L^2(\Omega)$.

Proposition 1.12. *The space $(\mathcal{L}^+, d)$ is complete. It is also separable if $(\Omega, \mathcal{B})$ is Borel.*

The next proposition shows that the metric d on the space $\mathcal{L}^+$ corresponds to convergence in probability on the space of stochastic integrals $\{I(f), f \in \mathcal{L}^+\}$.

Proposition 1.13. *For $f_n \in \mathcal{L}^+$ and $c_n \in \mathbb{R}$, we have that $I(f_n) + c_n$ converges to a random variable ξ in probability, as $n \to \infty$, if and only if, there exists some $f \in \mathcal{L}^+$ and $c \in \mathbb{R}$, such that $d(f_n - f) + |c_n - c| \to 0$, as $n \to \infty$. In this case, $\xi = I(f) + c$ a.s.*

The next theorem, which is analogous to Theorem 1.8, shows the uniqueness of the minimal spectral representation for i.d. processes.

Theorem 1.14. *Let $X = \{X(t), t \in T\}$ be an i.d. process. If $\{f_t^{(i)}, t \in T\}$ are two minimal spectral representations of X defined on σ–finite Borel spaces $(\Omega_i, \mathcal{B}_i, \mu_i)$, $i = 1, 2$, then there is a measure space isomorphism $\Phi : \Omega_1 \to \Omega_2$ (defined modulo null sets) such that for all $t \in T$,*

$$f_t^{(1)}(\omega) = f_t^{(2)} \circ \Phi(\omega) \quad \textit{for } \mu_1 - \textit{a.a. } \omega \in \Omega_1. \tag{12}$$

The measure space isomorphism Φ is unique modulo null sets.

1.4. **Flow representations for stationary i.d. and max–i.d. processes.** A measure–preserving $\mathbb{R}^d$–action (or flow) on a measure space $(\Omega, \mathcal{B}, \mu)$ is a family $\{T_t\}_{t\in\mathbb{R}^d}$ of measure space isomorphisms $T_t : \Omega \to \Omega$ such that $T_0 = \mathrm{id}$ μ–a.e. and for every $t, s \in \mathbb{R}^d$, $T_t \circ T_s = T_{t+s}$ μ–a.e. The action is called measurable if $(t, \omega) \mapsto T_t(\omega)$ is a measurable map from $\mathbb{R}^d \times \Omega$ to Ω, where the former space is endowed with the product of the Borel σ–algebra on $\mathbb{R}^d$ and $\mathcal{B}$.

The next statement combines both the i.d. and max-i.d. cases and shows that one can associate stationary processes with measure–preserving actions. The common theme is the uniqueness.

Theorem 1.15. *Let $X = \{X(t), t \in \mathbb{R}^d\}$ be a stationary, stochastically continuous, max–i.d. (resp., i.d., without Gaussian component) random field. Then there is a σ–finite Borel space $(\Omega, \mathcal{B}, \mu)$, a measurable, measure–preserving $\mathbb{R}^d$–action $\{T_t\}_{t\in\mathbb{R}^d}$ on $(\Omega, \mathcal{B}, \mu)$, and a function $f_0 \in \mathcal{L}^\vee(\Omega)$ (resp., $f_0 \in \mathcal{L}^+(\Omega)$) such that we have*

$$\{X(t), t \in \mathbb{R}^d\} \overset{d}{=} \left\{ \int_\Omega^{\vee/+} (f_0 \circ T_t) d\Pi_\mu + c^{\vee/+}, t \in \mathbb{R}^d \right\}, \tag{13}$$

where Π_μ is a Poisson random measure on $(\Omega, \mathcal{B})$ with intensity measure μ, with $c^\vee = 0$ and $c^+ \in \mathbb{R}$.

Proof. Stochastic continuity implies that X satisfies Condition S. Thus, let $\{f_t, t \in \mathbb{R}^d\}$ be a minimal spectral representation of X on $[0, 1]$ endowed with a σ–finite measure (Theorems 1.6 & 1.11). By stationarity, $\{f_{t+s}, t \in \mathbb{R}^d\}$ is also a minimal spectral representation of X for every $s \in \mathbb{R}^d$. By the uniqueness of the minimal spectral representation (Theorems 1.8 & 1.14), there is an automorphism T_s of the measure space $(\Omega, \mathcal{B}, \mu)$ such that for every $t \in \mathbb{R}^d$, $f_{s+t} = f_t \circ T_s$, μ–a.e. Let us show that for every $s_1, s_2 \in \mathbb{R}^d$, $T_{s_1+s_2} = T_{s_1} \circ T_{s_2}$, μ–a.e. Indeed, we have for every $t \in \mathbb{R}^d$,

$$f_t \circ T_{s_1+s_2} = f_{s_1+s_2+t} = f_t \circ (T_{s_1} \circ T_{s_2}), \quad \mu - \text{a.e.}$$

Since the automorphisms in the statements of Theorems 1.8 & 1.14 are unique, we have $T_{s_1+s_2} = T_{s_1} \circ T_{s_2}$, μ–a.e. From the definition of the spectral representation we obtain (13). In the sum–i.d. case note also that the constant c^+ appearing in (13) does not depend on $t \in \mathbb{R}^d$ by stationarity.

The flow $\{T_t\}_{t\in\mathbb{R}^d}$ constructed in this way need not in general be measurable. However, it was shown in [18] that each T_t can be modified on a set of μ–measure zero so that the flow becomes measurable. □

We call the pair $(f_0, \{T_t\}_{t\in\mathbb{R}^d})$ a flow representation of X on $(\Omega, \mathcal{B}, \mu)$. It follows from the proof of Theorem 1.15 that the flow representation can be chosen to be minimal meaning that the σ–algebra generated by $\{f_0 \circ T_t, t \in \mathbb{R}^d\}$ coincides with $\mathcal{B}$ up to μ–zero sets and the set of functions $\{f_0 \circ T_t,\ t \in \mathbb{R}^d\}$ has full support.

Theorem 1.16. *Let $X = \{X(t), t \in \mathbb{R}^d\}$ be a stationary, stochastically continuous max–i.d. or i.d. random field. The minimal flow representation of X is unique in the following sense. If $(f_0^{(i)}, \{T_t^{(i)}\}_{t\in\mathbb{R}^d})$ are two minimal flow representations of X on σ–finite Borel spaces $(\Omega_i, \mathcal{B}_i, \mu_i)$, $i = 1, 2$, then there is a measure space isomorphism $\Phi : \Omega_1 \to \Omega_2$ (defined modulo null sets) such that $f_0^{(1)} = f_0^{(2)} \circ \Phi$, μ_1–a.e., and for all $t \in \mathbb{R}^d$,*

$$\Phi \circ T_t^{(1)} = T_t^{(2)} \circ \Phi, \quad \mu_1 - a.e. \tag{14}$$

The isomorphism Φ is unique modulo null sets.

Proof. By assumption, $\{f_0^{(i)} \circ T_t^{(i)}, t \in \mathbb{R}^d\}$, $i = 1, 2$, are two minimal spectral representations of X. By the uniqueness of the minimal spectral representation, there is a measure space isomorphism $\Phi : \Omega_1 \to \Omega_2$ such that for every $t \in \mathbb{R}^d$,

$$(f_0^{(1)} \circ T_t^{(1)}) = (f_0^{(2)} \circ T_t^{(2)}) \circ \Phi, \quad \mu_1 - \text{a.e.} \tag{15}$$

Replacing t by $t + s$ and taking the composition of both sides with $T_{-s}^{(1)}$ from the right, we obtain

$$(f_0^{(1)} \circ T_t^{(1)}) = (f_0^{(2)} \circ T_t^{(2)}) \circ (T_s^{(2)} \circ \Phi \circ T_{-s}^{(1)}), \quad \mu_1 - \text{a.e.} \tag{16}$$

It follows from (15) and (16) that Φ and $T_s^{(2)} \circ \Phi \circ T_{-s}^{(1)}$ are two measure space isomorphisms each linking the representations $\{f_0^{(i)} \circ T_t^{(i)}, t \in \mathbb{R}^d\}$, $i = 1, 2$. By the last statement of Theorem 1.8 or Theorem 1.14 these isomorphisms should be equal up to μ_1–zero sets. This yields (14). □

2. Examples of max–i.d. processes

2.1. Max–stable processes. Max–stable processes form a subclass of the max–i.d. processes. Fix $\alpha > 0$. A process $X = \{X(t), t \in T\}$ is called (α–Fréchet) max–stable if for every $n \in \mathbb{N}$ the process $X_1 \vee \ldots \vee X_n$ has the same law as $n^{1/\alpha} X$, where $X_1, \ldots, X_n$ are i.i.d. copies of X. The marginal distributions of X are α–Fréchet distributions of the form $\mathbb{P}[X(t) \le x] = \exp\{-\sigma^\alpha(t) x^{-\alpha}\}$, $x > 0$. Here, $\sigma(t) > 0$ is called the scale parameter of $X(t)$.

For max–stable processes, a theory of spectral representations has been developed; see [4, 5, 33, 12, 37]. We will explain the connection to the max–i.d. spectral representations developed here. Let $L_+^\alpha(\Omega', \mathcal{B}', \mu')$ be the set of measurable functions $g : \Omega' \to [0, \infty)$ such that $\int_{\Omega'} g^\alpha d\mu' < \infty$. Let $\Omega = \Omega' \times (0, \infty)$ be equipped with the product σ–algebra $\mathcal{B}$ and with a measure $\mu = \mu' \times \alpha u^{-(\alpha+1)} du$. A collection of functions $\{g_t, t \in T\} \subset L_+^\alpha(\Omega', \mathcal{B}', \mu')$ is called a spectral representation of a max–stable process $\{X(t), t \in T\}$ if

$$\{X(t), t \in T\} \overset{d}{=} \left\{ \bigvee_{i \in \mathbb{N}} u_i g_t(\omega_i'), t \in T \right\}. \tag{17}$$

where $\{(\omega_i', u_i), i \in \mathbb{N}\}$, are points of the Poisson process Π_μ with intensity μ on Ω. The process X, being also max–i.d., must admit a spectral representation in the sense of Section 1.2. This representation can be constructed as follows. Define $f_t : \Omega \to [0, \infty)$ by $f_t(\omega', u) = u g_t(\omega')$, $t \in T$. Then, (17) implies that $\{f_t, t \in T\}$ is a spectral representation of X viewed as a max–i.d. process.

A spectral representation $\{g_t, t \in T\} \subset L_+^\alpha(\Omega', \mathcal{B}', \mu')$ of a max–stable process X is called minimal (see [12, 36, 37]) if (i) $\operatorname{supp}\{g_t,\ t \in T\} = \Omega' \mod \mu'$ and (ii) $\sigma\{g_t/g_s,\ t, s \in T\} = \mathcal{B}' \mod \mu'$.

Lemma 2.1. *In the above context, if $\{g_t, t \in T\}$ is a minimal spectral representation of a max–stable process X, then $\{f_t, t \in T\}$ is a minimal spectral representation of X as a max–i.d. process.*

Proof. Notice that $(f_t/f_s)(x, y) = (g_t/g_s)(y)$, (with $0/0$ is interpreted as 0) does not depend on x. Therefore, $\rho(F) := \sigma\{f_t/f_s,\ t, s \in T\} = \mathbb{R}_+ \times \sigma\{g_t/g_s,\ t, s \in T\}$, which is (mod μ) equivalent to $\mathbb{R}_+ \times \mathcal{B}_{\Omega'}$ by Condition (ii). We also have that g_t is (mod μ) measurable with respect to (w.r.t.) $\mathbb{R}_+ \times \mathcal{B}_{\Omega'}$ (mod μ) and since $\rho(F) = \mathbb{R}_+ \times \mathcal{B}_{\Omega'}$ (mod μ), it follows that g_t is (mod μ) measurable w.r.t. $\sigma(F) := \sigma\{f_t,\ t \in T\} (\supset \rho(F))$. Therefore, $(x, y) \mapsto x \mathbb{1}_{\{\operatorname{supp}(g_t)\}}(y) = f_t(x, y)/g_t(y)$ is (mod μ) measurable w.r.t. $\sigma(F)$. Now, the full support condition (i) implies also that $(x, y) \mapsto x$ is (mod μ) measurable w.r.t. $\sigma(F)$. This implies that $\mathcal{B}_{\mathbb{R}_+} \times \Omega'$ is included in $\sigma(F)$ (mod μ). Since also $\mathbb{R}_+ \times \mathcal{B}_{\Omega'}$ is (mod μ) included in $\sigma(F)$, it follows that $\mathcal{B}_{\mathbb{R}_+} \times \mathcal{B}_{\Omega'}$ is (mod μ) contained in $\sigma(F)$. This shows that $\mathcal{B}_{\mathbb{R}_+} \otimes \mathcal{B}_{\Omega'} \equiv \sigma(\mathcal{B}_{\mathbb{R}_+} \times \mathcal{B}_{\Omega'}) = \sigma(F)$ (mod μ). This shows that $\{f_t, t \in T\}$ is a minimal representation of X since Condition (i) for $\{g_t, t \in T\}$ implies also the full support condition for the f_t's. □

Remark 2.2. Lemma 2.1 shows that all previous results on max-stable processes that rely on the notion of minimality can be obtained via the new notion of minimality. The following construction due to Maharam gives the precise connection between the "old" and "new" spectral representations in the case of stationary processes.

Let $X = \{X(t), t \in \mathbb{R}^d\}$ be a stationary stochastically continuous max–stable process with α–Fréchet margins. Then by [5], there is a *non–singular* flow T'_t on a σ–finite Borel space $(\Omega', \mathcal{B}', \mu')$ and a function $g_0 \in L^\alpha_+(\Omega', \mathcal{B}', \mu')$ such that $\{g_t, t \in \mathbb{R}^d\}$ is a minimal spectral representation of X, where

$$g_t = \Big(\frac{d\mu' \circ T'_t}{d\mu'}\Big)^{1/\alpha} g_0 \circ T'_t, \quad t \in \mathbb{R}.$$

The process X, being max–stable, is also max–i.d. Let us construct the flow representation of X in the sense of Section 1.4. We shall employ the Maharam construction [19, 1]. Consider the mappings $T_t : \Omega \to \Omega$ defined by

$$T_t(u, \omega') := \left(\left(\frac{d\mu' \circ T'_t}{d\mu'}(\omega')\right)^{1/\alpha} u, T'_t(\omega')\right), \tag{18}$$

where $(t, \omega') \mapsto d(\mu' \circ T'_t)/d\mu'(\omega')$ is a measurable version of the Radon–Nikodym derivatives (see e.g. Theorem A.1 in [17]). It is easy to see that $\{T_t\}_{t\in\mathbb{R}}$ is a measurable flow, which is measure–preserving (see e.g. [19, 1]). Now, (18) implies that $(f_0, \{T_t\}_{t\in\mathbb{R}^d})$ is a flow representation of X in the sense of Section 1.4.

2.2. **Independent random variables.** Any collection $\{X(t), t \in T\}$ of independent random variables forms a max–i.d. process. To see this, take any $n \in \mathbb{N}$ and let $X_{i,n}(t), 1 \le i \le n, t \in T$, be independent random variables such that $\mathbb{P}[X_{i,n}(t) \le x] = (\mathbb{P}[X(t) \le x])^{1/n}$. Then, $\{X(t), t \in T\}$ has the same law as $\{\vee_{i=1}^n X_{i,n}(t), t \in T\}$, thus showing the max–i.d. property. Assume that T is countable. Then, Condition S is satisfied. The minimal spectral representation of $\{X(t), t \in T\}$ can be constructed as follows. As always, we assume that $\operatorname{essinf} X(t) = 0$ and, additionally, $\mathbb{P}[X(t) = 0] < 1$ for all $t \in T$. Let $\Omega = T \times (0, \infty)$ be endowed with the product of the power set 2^T and the Borel σ–algebra on $(0, \infty)$. Define a measure μ on Ω by $\mu(\{t\} \times [x, \infty)) = -\log \mathbb{P}[X(t) < x]$, $t \in T$, $x > 0$. In this way, Ω turns into a σ–finite Borel space. Define the functions $f_t : \Omega \to \mathbb{R}$, $t \in T$, by

$$f_t(s, x) = \begin{cases} x, & t = s, \\ 0, & t \neq s, \end{cases} \quad s \in T, \quad x > 0.$$

Then, $\{f_t, t \in T\}$ is a minimal spectral representation of $\{X_t, t \in T\}$. If $T = \mathbb{Z}$ and the random variables X_t are i.i.d., then X is stationary and we can define a (discrete time) flow representation by setting $T_t(s, x) = (s - t, x)$, $t \in \mathbb{Z}$, and noting that $f_t = f_0 \circ T_t$.

2.3. **Mixed moving maximum processes.** Let $\{U_i, i \in \mathbb{N}\}$ (interpreted as storm centers) be the points of a Poisson process on $\mathbb{R}^d$ with constant intensity λ. Let $\{F(t), t \in \mathbb{R}^d\}$ be a measurable random process with values in $[0, \infty)$ such that for every $a > 0$, the Lebesgue measure of the set $\{t \in \mathbb{R}^d : F(t) > a\}$ is finite a.s. Let F_n, $n \in \mathbb{N}$, be i.i.d. copies of F (storms), which are independent from the Poisson process $\{U_i, i \in \mathbb{N}\}$ of storm centers. Define a process $\{X(t), t \in \mathbb{R}^d\}$ by

$$X(t) = \sup_{i\in\mathbb{N}} F_i(t - U_i). \tag{19}$$

Then, it is readily verified (e.g., by the translation invariance and thinning properties of the point process $\{U_i, i \in \mathbb{N}\}$) that X is a stationary max–i.d. process, called a *mixed moving maximum process.*

The flow representation of X can be constructed as follows. Let ν be the probability law of F on the space $E := \mathbb{R}^{\mathbb{R}^d}$. Let Ω be the product space $E \times \mathbb{R}^d$ endowed with the measure $\mu := \nu \times \lambda ds$. Then, we may view $\Pi_\mu := \{(F_i, U_i), i \in \mathbb{N}\}$ as a Poisson point process on Ω with intensity μ. Define a flow $\{T_t\}_{t\in\mathbb{R}^d}$ on Ω and a function $f_0 : \Omega \to [0,\infty)$ by

$$T_t(\varphi, s) = (\varphi, s - t), \quad f_0(\varphi, s) = \varphi(-s), \quad (\varphi, s) \in \Omega,\ \varphi : \mathbb{R}^d \to \mathbb{R}.$$

This defines a flow representation of X. To see this, note that

$$\int_\Omega^\vee (f_0 \circ T_t) d\Pi_\mu = \sup_{i\in\mathbb{N}} (f_0 \circ T_t)(F_i, U_i) = \sup_{i\in\mathbb{N}} F_i(t - U_i), \quad t \in \mathbb{R}^d.$$

Of course, the space Ω is not Borel and the flow $\{T_t\}_{t\in\mathbb{R}^d}$ is not measurable. If, however, the process F is defined on a Borel space, we can use this space as E in the above construction to obtain a flow representation in the sense of Theorem 1.15.

As in the max–stable (or sum–stable) case [27] one obtains the following result.

Proposition 2.3. *A stationary max–i.d. process is of mixed moving maximum type iff the flow generating X is dissipative (see, e.g., [1] for a definition).*

2.4. **Max–i.d. processes associated to Poisson line processes.** Instead of taking points of a Poisson process as storm centers in (19), we can also take lines of a Poisson line process as storm centers. Let $\mathbb{T} = \mathbb{R}/(2\pi\mathbb{Z})$ be identified with the unit circle. Each point (r, φ) in $\mathbb{L} := \mathbb{R} \times \mathbb{T}$ corresponds to an oriented line in $\mathbb{R}^2$ which passes through the point $(r\cos\varphi, r\sin\varphi)$ in the direction of the vector $(-\sin\varphi, \cos\varphi)$. In this way, $\mathbb{L}$ can be identified with the set of all oriented lines in $\mathbb{R}^2$. Take a Poisson point process $\Pi = \{(r_i, \varphi_i), i \in \mathbb{N}\}$ on $\mathbb{L}$ whose intensity is $\lambda dr \times d\varphi$, where $\lambda > 0$ is constant. The corresponding random set of lines in $\mathbb{R}^2$ is called the Poisson line process and is interpreted as the set of storm centers. Its law is invariant with respect to translations of $\mathbb{R}^2$; see, e.g., [16]. Let $\{F(t), t \in \mathbb{R}\}$ be a process with values in $[0,\infty)$ such that for every $a > 0$, the Lebesgue measure of the set $\{t \in \mathbb{R} : F(t) > a\}$ is finite a.s. Let F_i, $i \in \mathbb{N}$, be i.i.d. copies of F (storms). Define a process $\{\eta(x, y), (x, y) \in \mathbb{R}^2\}$ by

$$\eta(x, y) = \max_{i\in\mathbb{N}} F_i(x\cos\varphi_i + y\sin\varphi_i - r_i). \tag{20}$$

Note that $|x\cos\varphi + y\sin\varphi - r|$ is the distance from the point (x, y) to the line corresponding to $(r, \varphi) \in \mathbb{L}$.

Proposition 2.4. *η is a stationary max–i.d. process on $\mathbb{R}^2$.*

One can construct also max–stable processes of this type. Fix $\alpha > 0$. Start with a Poisson process $\Phi = \{(r_i, \varphi_i, z_i), i \in \mathbb{N}\}$ on $\mathbb{L} \times (0,\infty)$ with intensity $\lambda dr \times d\varphi \times \alpha z^{-(\alpha+1)}dz$. Let $\{F(t), t \in \mathbb{R}\}$ be a process with values in $[0,\infty)$ such that $\mathbb{E}\int_\mathbb{R} F^\alpha(r)dr < \infty$. Let $\{F_i, i \in \mathbb{N}\}$ be independent copies of F. Define

$$\eta(x, y) = \max_{i\in\mathbb{N}} z_i F_i(x\cos\varphi_i + y\sin\varphi_i - r_i). \tag{21}$$

Proposition 2.5. *η is a stationary max–stable process on $\mathbb{R}^2$ with α–Fréchet margins.*

Max–stability follows directly from the properties of the Poisson processes and stationarity is the consequence of the stationarity of the Poisson line process. For simplicity, we considered processes based on Poisson line processes in $\mathbb{R}^2$, but a similar construction is possible for Poisson processes



Figure 1. The left and center panels show realizations of the stationary max–i.d. and max–stable random fields η in (20) and (21), respectively, with $h(t) = e^{-|t|}$. The right panel shows a realization of a non-ergodic min–i.d. Penrose–type random field generated by a stationary Gaussian process.

of k–dimensional affine subspaces in $\mathbb{R}^d$ for every $k < d$. For $k = 0$ we recover the mixed moving maxima processes, for $k \neq 0$ we conjecture that the processes do not have a mixed moving maximum representation. Rather, they should be generated by null–recurrent flows (see, e.g., Samorodnitsky [30], and also [28, 12, 37]).

2.5. **Penrose min–i.d. random fields.** The next family of examples generalizes the processes considered by Penrose [22, 23, 24]. Let $\Pi = \{U_i, i \in \mathbb{Z}\}$ be the points of Poisson process on $\mathbb{R}^k$ with a constant intensity λ. Let $\{\xi_i(t), t \in \mathbb{R}^d\}$, $i \in \mathbb{Z}$, be independent copies of a random field $\{\xi(t), t \in \mathbb{R}^d\}$ with values in $\mathbb{R}^k$ which has stationary increments. Let $|\cdot|$ be the Euclidean norm. Define

$$X(t) = \min_{i \in \mathbb{Z}} |U_i + \xi_i(t)|.$$

Proposition 2.6. *The process X is stationary, min–i.d. process (that is, $-X$ is max–i.d.).*

The min–i.d. property follows directly from the fact that for every $n \in \mathbb{N}$, we can represent Π as a union of n independent Poisson processes with constant intensity $\frac{\lambda}{n}$. The stationarity of X follows from the stationarity of increments of ξ; see Proposition 2.1 in [13]. To construct concrete families of examples one may take $k = 1$ and ξ to be the zero–mean Gaussian process defined on $\mathbb{R}^d$ with covariance function $\mathbb{E}[\xi(t)\xi(s)] = \frac{\sigma^2}{2}(|t|^{2H} + |s|^{2H} - |t-s|^{2H})$, where $H \in (0, 1]$ is the Hurst exponent and $\sigma^2 > 0$. Min–i.d. processes of this type appeared in [14] as limits of pointwise minima (in the sense of absolute value) of independent Gaussian processes. Also, one can take $d = 1$, $k \in \mathbb{N}$ arbitrary and let ξ be the $\mathbb{R}^k$–valued Brownian motion. We conjecture that the corresponding Penrose processes are of mixed moving maximum type for $k \geq 3$ (since the Brownian motion is transitive in this case) and are generated by null flows if $k \leq 2$ (see e.g., [30, 28, 12, 37]).

2.6. **Stationary union–i.d. random sets.** A measurable process $\{X(t), t \in \mathbb{R}^d\}$ taking only values $0, 1$ can be identified with the random set $S := \{t \in \mathbb{R}^d : X(t) = 1\}$. Note that we



Figure 2. Realizations of Penrose–type min–i.d. random fields driven by isotropic Lévy fractional Brownian motions defined on $\mathbb{R}^2$ with Hurst exponents $H = 0.1$, 0.5, and 0.9, respectively, left to right.

do not require the sets to be, say, closed. If the process X is max–i.d., then the corresponding random set S is *union–i.d.* and vice versa. This means that for every $n \in \mathbb{N}$ we can find i.i.d. random sets $C_1, \dots, C_n$ such that S has the same finite–dimensional distributions as $C_1 \cup \dots \cup C_n$; see [21], Chapter 4. The process X is stochastically continuous iff the random set S is stochastically continuous in the following sense: for every $t \in \mathbb{R}^d$, $\lim_{s\to t} \mathbb{P}[t \in S, s \notin S] = \lim_{s \to t} \mathbb{P}[t \notin S, s \in S] = 0$. Using Theorem 1.15 we can describe all stationary stochastically continuous union–i.d. random sets. It turns out that any such set has the following structure. Take a Poisson process on a σ–finite measure space and let A be a subset of finite measure of this space. Shift all points of the Poisson process by some measure–preserving $\mathbb{R}^d$–flow and look at those moments of time at which A contains at least one point of the Poisson process. These moments of time form a union–i.d. random set in $\mathbb{R}^d$, and vice versa, any union–i.d. set is equal (in the sense of finite–dimensional distributions) to a set of this form.

Theorem 2.7. *Let S be a stationary, stochastically continuous, union–i.d. random set in $\mathbb{R}^d$. Then there is a σ–finite Borel space $(\Omega, \mathcal{B}, \mu)$, a measurable, measure–preserving $\mathbb{R}^d$–action $\{T_t\}_{t\in\mathbb{R}^d}$ on $(\Omega, \mathcal{B}, \mu)$, and a set $A \in \mathcal{B}$ with $\mu(A) < \infty$ such that*

$$S \overset{d}{=} \left\{ t \in \mathbb{R}^d : \Pi_\mu(T_t^{-1}(A)) \neq 0 \right\}, \tag{22}$$

where Π_μ is a Poisson random measure on $(\Omega, \mathcal{B})$ with intensity measure μ.

Proof. Let $\{X(t), t \in \mathbb{R}^d\}$ be the $\{0, 1\}$–valued max–i.d. process corresponding to S, that is, $X(t) = \mathbb{1}_{t\in S}$. Then, X has a flow representation in the sense of Theorem 1.15. The function f_0 in this representation takes only values $0, 1$, (mod μ). That is, $f = \mathbb{1}_A$ for some set $A \in \mathcal{B}$. Note that A has finite measure since $f \in \mathcal{L}^\vee$. Since $f_0 \circ T_t = \mathbb{1}_{T_t^{-1}A}$, the statement of the theorem follows. □

For example, let Ω be the space $\mathbb{R}^d$ endowed with the Lebesgue measure. Consider a flow $T_t(\omega) = \omega - t$, $\omega, t \in \mathbb{R}^d$. Let $A \subset \mathbb{R}^d$ be a Borel set of finite measure and let $\Pi = \{U_i, i \in \mathbb{N}\}$ be a unit intensity Poisson process on $\mathbb{R}^d$. Then, the corresponding union–i.d. random set S has the form $S = \cup_{i\in\mathbb{N}}(U_i - A)$ and is known in the literature as the Boolean model with (non-random) grain A. One can also take the space Ω to be a product of some probability space E and the space $\mathbb{R}^d$ endowed with the Lebesgue measure, and define a flow T_t on Ω by shifting the second coordinate. In this way one obtains a Boolean model with random grains. This construction is a special case of mixed moving maximum processes.

3. Proofs

3.1. **Lemma on conjugacy between collections of functions.**

The following lemma (which is of independent interest) is used in the proofs of Theorems 1.8 and 1.14. It states that any two families of equimeasurable functions on σ–finite Borel spaces can be related to each other by a measure space isomorphism.

Lemma 3.1. *Let $(\Omega_i, \mathcal{B}_i)$ be two Borel spaces endowed with measures $\mu_i,\ i = 1, 2$. Consider two families of measurable functions $f_t^{(i)} : \Omega_i \to \mathbb{R},\ t \in T,\ i = 1, 2$, and define two measurable mappings $F_i : (\Omega_i, \mathcal{B}_i) \to (\mathbb{R}^T, \mathcal{B})$ by $F_i(\omega) = (f_t^{(i)}(\omega))_{t\in T},\ \omega \in \Omega_i,\ i = 1, 2$. Here, $\mathcal{B}$ is the product σ–algebra on $\mathbb{R}^T$. Assume that*

1. *$\sigma\{f_t^{(i)},\ t \in T\} = \mathcal{B}_i \mod \mu_i,\ i = 1, 2$.*
2. *The families $\{f_t^{(1)}, t \in T\}$ and $\{f_t^{(2)}, t \in T\}$ are equimeasurable, that is, the induced measures $\mu_1 \circ F_1^{-1}$ and $\mu_2 \circ F_2^{-1}$ are equal on $(\mathbb{R}^T, \mathcal{B})$.*

Then, there exists a measure space isomorphism $\Phi : \Omega_1 \to \Omega_2$ (defined modulo null sets) such that for all $t \in T$, we have $f_t^{(1)} = f_t^{(2)} \circ \Phi$, μ_1–a.e. Moreover, Φ with this property is unique modulo null sets.

Proof. It will be convenient to identify the sets in $\mathcal{B}_i$ that are equal modulo $\mu_i,\ i = 1, 2$. Formally, let $\mathcal{I}_i \subset \mathcal{B}_i$ be the σ–ideals of μ_i–null sets in the spaces $(\Omega_i, \mathcal{B}_i, \mu_i)$ (see e.g. Ch. II.21 in [32]) and let $[\mathcal{B}_i] := \mathcal{B}_i/\mathcal{I}_i$ be the corresponding factor σ–fields, $i = 1, 2$. The elements of $[\mathcal{B}_i]$ are the equivalence classes $[B] = \{A \in \mathcal{B}_i : \mu_i(A\Delta B) = 0\}$, where $B \in \mathcal{B}_i,\ i = 1, 2$.

We shall define next a σ–isomorphism $U : [\mathcal{B}_1] \to [\mathcal{B}_2]$, i.e. a bijective mapping that preserves countable unions and complements. For all $B \in \mathcal{B}_1$, we set

$$U([B]) := [F_2^{-1}(A)],\ \text{where}\ [F_1^{-1}(A)] = [B]. \tag{23}$$

Note that such an $A \in \mathcal{B}$ exists since by assumption $F_1^{-1}(\mathcal{B}) = \sigma\{f_t^{(1)},\ t \in T\} = \mathcal{B}_1 \mod \mu_1$. One can readily see that the mapping U is a well-defined σ–isomorphism. Indeed, since $\mu_1 \circ F_1^{-1} = \mu_2 \circ F_2^{-1}$, for every $A', A'' \in \mathcal{B}$,

$$\mu_1(F_1^{-1}(A')\Delta F_1^{-1}(A'')) = \mu_1(F_1^{-1}(A'\Delta A'')) = \mu_2(F_2^{-1}(A'\Delta A'')) = \mu_2(F_2^{-1}(A')\Delta F_2^{-1}(A'')).$$

Thus, $F_1^{-1}(A') = F_1^{-1}(A'') \mod \mu_1$, if and only if $F_2^{-1}(A') = F_2^{-1}(A'') \mod \mu_2$, and the definition of U does not depend on the choice of the representative B of the equivalence class $[B]$ and on the choice of A in (23). This shows, moreover, that $[B'] = [B'']$ if and only if $U([B']) = U([B''])$, that is, U is *injective*. On the other hand, since $F_2^{-1}(\mathcal{B}) = \sigma\{f_t^{(2)},\ t \in T\} = \mathcal{B}_2 \mod \mu_2$, for all $B \in \mathcal{B}_2$, we have $[F_2^{-1}(A)] = [B]$, for some $A \in \mathcal{B}$ and hence $U([F_1^{-1}(A)]) = [B]$. This shows that U is *onto* and hence a bijection. Also, since $\mu_1(F_1^{-1}(A)) = \mu_2(F_2^{-1}(A))$, we have by (23) that U is measure–preserving. Since U clearly preserves the countable unions and complements, it is a σ–isomorphism.

Since $(\Omega_i, \mathcal{B}_i)$ are Borel spaces, Theorem 32.5 of [32] implies that the σ–isomorphism U is induced by some measure space isomorphism $\Phi : \Omega_1 \to \Omega_2$ in the following sense:

$$U([B]) = [\Phi(B)], \quad B \in \mathcal{B}_1. \tag{24}$$

Let us fix some $t \in T$ and show that $f_t^{(1)} = f_t^{(2)} \circ \Phi$ holds μ_1–a.e. Let I be a Borel subset of $\mathbb{R}$ and consider the cylinder set $A = \{\varphi : T \to \mathbb{R} : \varphi(t) \in I\} \subset \mathbb{R}^T$. We have

$$\begin{aligned}[(f_t^{(2)} \circ \Phi)^{-1}(I)] &= [\Phi^{-1}((f_t^{(2)})^{-1}(I))] = U^{-1}([(f_t^{(2)})^{-1}(I)]) \\ &= U^{-1}([F_2^{-1}(A)]) = [F_1^{-1}(A)] = [(f_t^{(1)})^{-1}(I)].\end{aligned} \tag{25}$$

Assume that $f_t^{(1)} \neq f_t^{(2)} \circ \Phi$ on $D \in \mathcal{B}_1$ with $\mu_1(D) > 0$. Then we can find an $\varepsilon > 0$ and a measurable set $D' \subset D$ with $\mu_1(D) > 0$ such that $|f_t^{(1)} - f_t^{(2)} \circ \Phi| > \varepsilon$ everywhere on D'. Further, we can find a $k \in \mathbb{Z}$ and a measurable set $D'' \subset D'$ with $\mu_1(D'') > 0$ such that with $I = [k\varepsilon, (k+1)\varepsilon)$, we have $f_t^{(1)} \in I$ everywhere on D''. It then follows that $f_t^{(2)} \circ \Phi \notin I$ on D''. But this contradicts (25), which implies $f_t^{(1)} = f_t^{(2)} \circ \Phi$, μ_1–a.e.

To complete the proof of the lemma, we need to show that the isomorphism Φ is unique modulo null sets. Assume that $\widetilde{\Phi} : \Omega_1 \to \Omega_2$ is another measure space isomorphism (defined modulo null sets) such that for all $t \in T$, $f_t^{(1)} = f_t^{(2)} \circ \widetilde{\Phi}$, μ_1–a.e. Let $\widetilde{U} : [\mathcal{B}_1] \to [\mathcal{B}_2]$ be the induced factor σ–algebra isomorphism $\widetilde{U}([A]) := [\widetilde{\Phi}(A)]$, $A \in \mathcal{B}_1$. Then, for every Borel set $I \subset \mathbb{R}$ and every $t \in T$, we have

$$\begin{aligned}\widetilde{U}([(f_t^{(1)})^{-1}(I)]) &= [\widetilde{\Phi}((f_t^{(1)})^{-1}(I))] = [(f_t^{(2)})^{-1}(I)] \\ &= [\Phi((f_t^{(1)})^{-1}(I))] = U([(f_t^{(1)})^{-1}(I)]).\end{aligned}$$

Since sets of the form $(f_t^{(1)})^{-1}(I)$, where $I \subset \mathbb{R}$ is Borel, generate the σ–algebra $\mathcal{B}_1 \mod \mu_1$ and since $U, \ \widetilde{U} : [\mathcal{B}_1] \to [\mathcal{B}_2]$ are σ–isomorphisms, it follows that $U([B]) = \widetilde{U}([B])$ for every $B \in \mathcal{B}_1$.

Let now $\Psi := \Phi^{-1} \circ \widetilde{\Phi} : \Omega_1 \to \Omega_1$. Clearly, Ψ is a measure space isomorphism (defined modulo null sets), and as shown above, $[\Psi(A)] = [A]$ for all $A \in \mathcal{B}_1$. We will use the fact that $(\Omega_1, \mathcal{B}_1)$ is a Borel space to show that $\Psi = \mathrm{id} \mod \mu_1$, which will complete the proof of the result.

By Kuratowski's Theorem, $(\Omega_1, \mathcal{B}_1)$ is isomorphic to either $(E, 2^E)$, where E is an at most countable set, or $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ – the real line equipped with the Borel σ–algebra. The discrete case is trivial. Suppose now the latter is true and without loss of generality let $(\Omega_1, \mathcal{B}_1) \equiv (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$. Let $\varepsilon > 0$ be arbitrary and suppose that $\mu_1(\{|\Psi - \mathrm{id}| > \varepsilon\}) > 0$, then for some $k \in \mathbb{Z}$, we have for $D := \{|\Psi - \mathrm{id}| > \varepsilon\} \cap [k\varepsilon, (k+1)\varepsilon)$ that $\mu_1(D) > 0$. But then $\Psi(x) \notin [k\varepsilon, (k+1)\varepsilon)$, for all $x \in D$, and hence $\Psi(D) \cap D = \emptyset$. This contradicts the fact that $[\Psi(D)] = [D]$ because $\mu_1(D) > 0$. Since $\varepsilon > 0$ was arbitrary, it follows that $\Psi = \mathrm{id} \mod \mu_1$ and the proof is complete. □

3.2. **Proofs in the max–i.d. case.**

Proof of Theorem 1.6. Write $\mathbb{R}_+ = [0, \infty)$. Let T_0 be the at most countable set appearing in Condition S. Let $\mathbb{R}_+^{T_0}$ be the space of functions $\varphi : T_0 \to \mathbb{R}_+$ endowed with the product σ–algebra $\mathcal{B}$. Denote by ν the exponent measure of the process $\{X(t), t \in T_0\}$; see Vatan [34]. It is a σ–finite measure on $\mathbb{R}_+^{T_0}$ such that for every $t_1, \dots, t_n \in T_0$ and $x_1, \dots, x_n > 0$ we have

$$\mathbb{P}\{X(t_j) < x_j, \ 1 \le j \le n\} = \exp\Big\{ -\nu\Big(\cup_{j=1}^n \{\varphi \in \mathbb{R}_+^{T_0} : \varphi(t_j) \ge x_j\}\Big)\Big\}. \tag{26}$$

We agree that $\nu(\{0\}) = 0$ (which is different from [34]). Taking the coordinate mappings $f_t : \mathbb{R}_+^{T_0} \to \mathbb{R}$, $f_t(\varphi) = \varphi(t)$, $t \in T_0$, we therefore obtain a spectral representation of $\{X(t), t \in T_0\}$ on $(\mathbb{R}_+^{T_0}, \mathcal{B})$. To see this, compare (3) and (26). Let $t \in T$ be arbitrary. Condition S states that there exists a sequence $\{t_n\}_{n\in\mathbb{N}} \subset T_0$ such that $X(t_n) \to X(t)$ in probability. Thus, the sequence $X(t_n)$ is Cauchy in probability. By the equality of the finite–dimensional distributions, the sequence $I(f_{t_n})$ is Cauchy in probability, and therefore, it converges in probability. By Theorem 4.5 in [2], there is a function $f_t \in \mathcal{L}^\vee(\mathbb{R}_+^{T_0}, \mathcal{B}, \nu)$ such that $I(f_{t_n})$ converges in probability to $I(f_t)$. Theorem 4.4 of [2] implies that the finite–dimensional distributions of $\{I(f_t),\ t \in T\}$ and $\{X(t),\ t \in T\}$ are equal, that is, the collection $\{f_t, t \in T\}$ is a spectral representation of $\{X(t), t \in T\}$ on $(\mathbb{R}_+^{T_0}, \mathcal{B})$. Since the coordinate functions f_t, $t \in T_0$, generate the product σ–algebra $\mathcal{B}$, and $\nu(\cap_{t\in T_0}\{f_t = 0\}) = \nu(\{0\}) = 0$, this representation is minimal. To complete the proof note that by Kuratowski's theorem, for at most countable T_0, the measurable space $(\mathbb{R}_+^{T_0}, \mathcal{B})$ is isomorphic to $[0, 1]$ endowed with the Borel σ–algebra. □

Proof of Theorem 1.8. As in Lemma 3.1 we define two measurable mappings $F_i : (\Omega_i, \mathcal{B}_i) \to (\mathbb{R}^T, \mathcal{B})$ by

$$F_i(\omega) = (f_t^{(i)}(\omega))_{t\in T}, \quad \omega \in \Omega_i, \quad i = 1, 2.$$

The first condition of Lemma 3.1 is satisfied by the assumption of minimality. We will show that the induced measures $\mu_1 \circ F_1^{-1}$ and $\mu_2 \circ F_2^{-1}$ are equal on $(\mathbb{R}^T, \mathcal{B})$. We will prove that for all $t_1, \dots, t_n \in T$ and all intervals $[x_1, y_1), \dots, [x_n, y_n) \subset \mathbb{R}$ we have

$$\mu_1\left(\cap_{j=1}^n \{x_j \le f_{t_j}^{(1)} < y_j\}\right) = \mu_2\left(\cap_{j=1}^n \{x_j \le f_{t_j}^{(2)} < y_j\}\right). \tag{27}$$

Recall that $\{f_t^{(i)}, t \in T\}$, $i = 1, 2$, are spectral representations of the same process X. By (3) we have that for all $x_1, \dots, x_n > 0$,

$$\mu_1\left(\cup_{j=1}^n \{f_{t_j}^{(1)} \ge x_j\}\right) = \mu_2\left(\cup_{j=1}^n \{f_{t_j}^{(2)} \ge x_j\}\right). \tag{28}$$

Note that $\mu_i(\{f_t^{(i)} > x\}) < \infty$ for all $x > 0$, $t \in T$, since $f_t^{(i)} \in \mathcal{L}^\vee$. Using this fact and the inclusion–exclusion formula we obtain that Relation (28) is also valid with the unions therein replaced by intersections. This proves that (27) holds provided that $0 < x_j < y_j$ for all $j = 1, \dots, n$. Note that this argument breaks down if $x_j = 0$ for some j since we cannot apply the inclusion–exclusion formula to sets of infinite measure. To show that the measures $\mu_1 \circ F_1^{-1}$ and $\mu_2 \circ F_2^{-1}$ agree on the "boundary" of $\mathbb{R}_+^T$ we need a separate argument.

We now show that (27) continues to hold even if some of the x_j's are allowed to be zero. We don't need to consider the case of negative x_j's since $f_t^{(i)} \ge 0$, μ_i–a.e., by definition of $\mathcal{L}^\vee$. By letting some of the x_j's go to 0 and using continuity of measure we obtain that (27) continues to hold if some of the sets of the form $\{x_j \le f_{t_j}^{(i)} < y_j\}$ therein are replaced by $\{0 < f_{t_j}^{(i)} < y_j\}$. By additivity of measure, the proof of (27) in full generality will be completed if we show that (27) continues to hold if some of the sets of the form $\{x_j \le f_{t_j}^{(i)} < y_j\}$ therein are replaced by $\{f_{t_j}^{(i)} = 0\}$. Let us make this statement precise. Take $l, m \in \mathbb{N}_0$, $s_1, \dots, s_l \in T$, $r_1, \dots, r_m \in T$ and $0 < u_1 < v_1, \dots, 0 < u_m < v_m$. Define two measurable sets $C_i \subset \Omega_i$, $i = 1, 2$, by

$$C_i = A_i \cap B_i, \quad A_i = \cap_{k=1}^l \{f_{s_k}^{(i)} = 0\}, \quad B_i = \cap_{j=1}^m \{u_j \le f_{r_j}^{(i)} < v_j\}, \quad i = 1, 2.$$

We will show that $\mu_1(C_1) = \mu_2(C_2)$. Suppose first that $m \neq 0$. Then, $\mu_i(C_i) = \mu_i(B_i) - \mu_i(B_i \cap D_i)$, where

$$D_i = \cup_{k=1}^{l} \{f_{s_k}^{(i)} > 0\} = \cup_{n\in\mathbb{N}} D_{i,n}, \quad D_{i,n} = \cup_{k=1}^{l} \left\{\frac{1}{n} \leq f_{s_k}^{(i)} < n\right\}, \quad i = 1, 2.$$

We have already shown that (27) holds if $x_j > 0$ for all $j = 1, \dots, n$. This implies that $\mu_1(B_1) = \mu_2(B_2)$ (where both terms are finite since $m \neq 0$). Also, by the inclusion–exclusion formula, $\mu_1(B_1 \cap D_{1,n}) = \mu_2(B_2 \cap D_{2,n})$ for every $n \in \mathbb{N}$. Note that $D_{i,1} \subset D_{i,2} \subset \dots$. Letting $n \to \infty$ and using the continuity of measure, we obtain $\mu(B_1 \cap D_1) = \mu_2(B_2 \cap D_2)$. This proves that $\mu_1(C_1) = \mu_2(C_2)$ in the case $m \neq 0$.

Consider now the case $m = 0$. In this case it is possible that $\mu_i(B_i) = \infty$ and the above argument breaks down. We show that $\mu_1(C_1) = \mu_2(C_2)$, or, equivalently, $\mu_1(A_1) = \mu_2(A_2)$. We will use the minimality and an exhaustion argument (cf. Lemma 1.0.7 in [1]) to show that there is a sequence $q_1, q_2, \dots \in T$ such that

$$\mu_i\left(\cap_{n\in\mathbb{N}} \{f_{q_n}^{(i)} = 0\}\right) = 0, \quad i = 1, 2. \tag{29}$$

Fix $i \in \{1, 2\}$. Since the measure μ_i is σ–finite, we can represent Ω_i as a disjoint union of sets $E_1, E_2, \dots \in \mathcal{B}_i$ such that $\mu_i(E_k) < \infty$, $k \in \mathbb{N}$. Let $e_k = \inf_Q \mu_i(\cap_{q\in Q}\{f_q^{(i)} = 0\} \cap E_k)$, where the infimum is taken over all at most countable sets $Q \subset T$. Clearly, $e_k < \infty$. For every $n \in \mathbb{N}$ we can find at most countable $Q_{kn} \subset T$ such that $\mu_i(\cap_{q\in Q_{kn}}\{f_q^{(i)} = 0\} \cap E_k) < e_k + \frac{1}{n}$. Since $Q_k := \cup_{n\in\mathbb{N}} Q_{kn}$ is at most countable, we have $e_k = \mu_i(F_k)$, where $F_k = \cap_{q\in Q_k}\{f_q^{(i)} = 0\} \cap E_k$. It follows that for every $t \in T$, $f_t^{(i)} = 0$ a.e. on F_k. Otherwise, we could consider $Q_k \cup \{t\}$ and arrive at a contradiction. By the assumption of minimality this implies that we must have $e_k = 0$. This holds for every $k \in \mathbb{N}$. The proof of (29) is completed by taking the union of the collections Q_k, $k \in \mathbb{N}$.

Consider measurable sets

$$G_{i,p} = A_i \cap \left(\cap_{k=1}^{p-1}\{f_{q_k}^{(i)} = 0\}\right) \cap \{f_{q_p}^{(i)} > 0\}, \quad p \in \mathbb{N}, \quad i = 1, 2.$$

We have $\mu_1(G_{1,p}) = \mu_2(G_{2,p})$ for every $p \in \mathbb{N}$. Indeed, by continuity of measure,

$$\mu_i(G_{i,p}) = \lim_{n\to\infty} \mu_i\left(A_i \cap \left(\cap_{k=1}^{p-1}\{f_{q_k}^{(i)} = 0\}\right) \cap \left\{\frac{1}{n} \leq f_{q_p}^{(i)} < n\right\}\right).$$

The right–hand side does not depend on $i = 1, 2$ as a particular case of $\mu_1(C_1) = \mu_2(C_2)$ in the case $m > 0$. It follows from (29) that

$$\mu_1(A_1) = \sum_{p=1}^{\infty} \mu_1(G_{1,p}) = \sum_{p=1}^{\infty} \mu_2(G_{2,p}) = \mu_2(A_2).$$

This completes the proof of (27).

It follows now from (27) that the measures $\mu_1 \circ F_1^{-1}$ and $\mu_2 \circ F_2^{-1}$ coincide on the semi–ring $\mathcal{C}$ consisting of sets of the form $\cap_{j=1}^{n}\{\varphi : T \to \mathbb{R} : x_j \leq \varphi(t_j) < y_j\}$, where $t_1, \dots, t_n \in T$, $[x_1, y_1), \dots, [x_n, y_n) \subset \mathbb{R}$. Note that $\mathcal{C}$ generates the product σ–algebra $\mathcal{B}$. Also, by (29), we can represent $\mathbb{R}^T$ as

$$\mathbb{R}^T = \cup_{n=1}^{\infty} \cup_{k=1}^{\infty} \left\{\varphi : T \to \mathbb{R} : k^{-1} \leq \varphi(q_n) < k\right\} \quad \text{mod } \mu_1 \circ F_1^{-1} \text{ and } \mu_2 \circ F_2^{-1}.$$

Note that the sets in the union in the right–hand side have finite $\mu_1 \circ F_1^{-1}$ (and $\mu_2 \circ F_2^{-1}$) measure and belong to the semiring $\mathcal{C}$. The uniqueness of the extension of measure theorem yields that $\mu_1 \circ F_1^{-1} = \mu_2 \circ F_2^{-1}$. The assumptions of Lemma 3.1 are verified. Lemma 3.1 yields (5) and completes the proof of the theorem. ☐

3.3. **Proofs in the i.d. case.**

We start with discussing some properties of the spectral representation. Note first that the functional I is not additive. Nevertheless, by (9) it follows that for all $f, g \in \mathcal{L}^+$

$$I(f) + I(g) = I(f+g) + \gamma(f,g), \quad \text{where } \gamma(f,g) := \int_\Omega \Big(a(f) + a(g) - a(f+g)\Big) d\mu. \tag{30}$$

The next result shows that γ in (30) is well defined and that this constant correction term can be controlled in terms of the metric d.

Lemma 3.2. *For all $f, g \in \mathcal{L}^+$, we have that $\int_\Omega |a(f) + a(g) - a(f+g)| d\mu < \infty$. Moreover, for γ and d as in* (30) *and* (11)*, we have*

$$|\gamma(f,g)| \le 3d(f+g)^2 + 2(d(f) + d(g))d(f+g).$$

Proof. Consider the integral defining $\gamma(f,g)$ over the sets $A := \{|f+g| > 1\}$, $B := A^c \cap \{|f| \le 1\} \cap \{|g| \le 1\}$ and $C := A^c \cap (\{|f| > 1\} \cup \{|g| > 1\})$, which form a disjoint partition of Ω.

Observe that over B the integrand is zero, since $a(f) = f$, $a(g) = g$ and $a(f+g) = f+g$ whenever $|f| \le 1$, $|g| \le 1$, and $|f+g| \le 1$. Note, on the other hand that the set $B^c = A \cup C \subset \{|f+g| > 1\} \cup \{|f| > 1\} \cup \{|g| > 1\}$ has a finite μ measure because f, g and $f+g$ belong to $\mathcal{L}^+$. Since $|a(f) + a(g) - a(f+g)| \le 3$ it therefore follows that $\int_\Omega |a(f) + a(g) - a(f+g)| d\mu < \infty$ and $\gamma(f,g)$ is well defined.

Over A, we have that

$$\int_{\{|f+g|>1\}} |a(f) + a(g) - a(f+g)| d\mu \le 3\mu\{|f+g| > 1\} \le 3d(f+g)^2.$$

Now, focus on the set C. The function a in (8) is Lipschitz and in fact $|a(x) + a(y)| \le |x+y|$ for all $x, y \in \mathbb{R}$. Therefore, $|a(f) + a(g) - a(f+g)| \le 2|f+g|$, and hence

$$\begin{aligned}\int_C |a(f) + a(g) - a(f+g)| d\mu &\le 2\int_{\{|f+g|\le 1\}} |f+g|(\mathbb{1}_{\{|f|>1\}} + \mathbb{1}_{\{|g|>1\}}) d\mu \\ &\le 2d(f+g)(d(f) + d(g)),\end{aligned}$$

where the last relation follows from the Cauchy–Schwartz inequality and the fact that $\mu\{|f| > 1\} \le d(f)^2$. Combining the above two bounds, we obtain the desired inequality. ☐

Relation (30) readily implies the following result on the sum of two spectral representations over the same space.

Proposition 3.3. *Consider two i.d. processes $X_t^{(i)} := I(f_t^{(i)}) + c_t^{(i)}$, $t \in T$, where $\{f_t^{(i)}\}_{t\in T} \subset \mathcal{L}^+(\Omega, \mathcal{B}, \mu)$, and $c_t^{(i)} \in \mathbb{R}$, $i = 1, 2$. Then, their sum has the following spectral representation:*

$$\{X_t^{(1)} + X_t^{(2)}\}_{t\in T} \overset{d}{=} \Big\{I(f_t^{(1)} + f_t^{(2)}) + c_t^{(1)} + c_t^{(2)} + \gamma(f_t^{(1)}, f_t^{(2)}), t \in T\Big\}.$$

Recall now that convergence in probability is metrized by the Ky Fan distance which is given by

$$d_{KF}(\xi,\eta) \equiv d_{KF}(\xi-\eta) := \inf\{\delta>0 \,:\, \mathbb{P}\{|\xi-\eta|\geq\delta\}\leq\delta\}. \tag{31}$$

The next proposition shows that the metric d on the space of integrands is comparable to the metric d_{KF} on the space of integrals.

Proposition 3.4. *For all $f\in\mathcal{L}^+$, we have*

$$d_{KF}(I(f)) \leq 2d(f)^{2/3} \quad \text{and} \quad 1-e^{-cd(f)^2} \leq 2d_{KF}(I(f)-I(f)') \leq 4d_{KF}(I(f)), \tag{32}$$

with $c=1-\sin(1)$, where $I(f)'$ is an independent copy of $I(f)$.

The following elementary inequality is used in the proof of Proposition 3.4.

Lemma 3.5. *Let X be a symmetric random variable. Then*

$$\sup_{|\theta|\leq1}(1-\mathbb{E}e^{i\theta X}) \leq 2d_{KF}(X). \tag{33}$$

Proof of Lemma 3.5. Since X is symmetric, we have that its characteristic function $\phi_X(\theta)=\mathbb{E}e^{i\theta X}$, $\theta\in\mathbb{R}$ is real and

$$(1-\phi_X(\theta)) = \int_{-\infty}^{\infty}(1-\cos(\theta x))F_X(dx).$$

Note that $0\leq 1-\cos(u)\leq u^2/2$, for all $u\in\mathbb{R}$. Thus, with $\varepsilon\in(0,1]$, we have

$$(1-\phi_X(\theta)) \leq \frac{|\theta\varepsilon|^2}{2}\int_{-\varepsilon}^{\varepsilon}F_X(dx) + \int_{|x|\geq\varepsilon}F_X(dx) \leq \varepsilon + \mathbb{P}\{|X|\geq\varepsilon\}$$

for all $|\theta|\leq 1\leq\sqrt{2/\varepsilon}$. The inequality (33) follows from the definition (31) of the Ky Fan distance functional. ☐

Proof of Proposition 3.4. We first prove the second inequality in (32). Let $X:=I(f)-I(f)'$, where $I(f)'$ is an independent copy of $I(f)$. Thus, in view of (9), X is symmetric with characteristic function

$$\phi_X(\theta) = |\mathbb{E}e^{i\theta I(f)}|^2 = \exp\Big\{-2\int_\Omega(1-\cos(\theta f))d\mu\Big\}, \quad \theta\in\mathbb{R}. \tag{34}$$

Now, by Lemma 3.5, we obtain

$$0\leq \sup_{|\theta|\leq1}(1-\phi_X(\theta)) \leq 2d_{KF}(X).$$

Thus, in view of (34), using the fact that the function $u\mapsto 1-e^{-2u}$, $u\geq0$ is strictly increasing, the above supremum can be taken inside the exponential, and hence

$$1-e^{-2A} := 1-\exp\Big\{-2\sup_{|\theta|\leq1}\int_\Omega(1-\cos(\theta f))d\mu\Big\} \leq 2d_{KF}(X). \tag{35}$$

We will focus on the term A above and obtain a lower bound for it. Notice that

$$\sup_{|\theta|\leq1}\int_{\{|f|\leq1\}}(1-\cos(\theta f))d\mu + \sup_{|\theta|\leq1}\int_{\{|f|>1\}}(1-\cos(\theta f))d\mu \leq A+A\equiv 2A.$$

Since $x^2/3 \le 1-\cos(x),\ |x|\le 1$, for the first term above, we have

$$\frac{1}{3}\int_{\{|f|\le 1\}} |f|^2 d\mu = \sup_{|\theta|\le 1} \frac{\theta^2}{3}\int_{\{|f|\le 1\}} |f|^2 d\mu \le \sup_{|\theta|\le 1}\int_{\{|f|\le 1\}} (1-\cos(\theta f))d\mu.$$

On the other hand, over the set $\{|f|>1\}$, we apply the inequality $\sup_{|\theta|\le 1}(1-\cos(\theta f)) \ge \int_0^1 (1-\cos(\theta f))d\theta = 1-\sin(f)/f$. By combining these two lower bounds, we obtain

$$\frac{1}{3}\int_{\{|f|\le 1\}} |f|^2 d\mu + \int_{\{|f|>1\}} \Big(1-\frac{\sin(f)}{f}\Big)d\mu \le 2A. \tag{36}$$

Also, since $1-\sin(x)/x \ge 1-\sin(1) =: c \approx 0.1585 > 0$, for all $|x|\ge 1$, we obtain further that

$$cd(f)^2 \le \frac{1}{3}\int_{\{|f|\le 1\}} |f|^2 d\mu + \int_{\{|f|>1\}} \Big(1-\frac{\sin(f)}{f}\Big)d\mu.$$

In view of (35), (36), and the monotonicity of $u \mapsto 1-e^{-u}$, we obtain $1-e^{-cd(f)^2} \le 2d_{KF}(X)$, which, since $d_{KF}(X) \equiv d_{KF}(I(f)-I(f)') \le 2d_{KF}(I(f))$, yields the second inequality in (32).

We now establish the first inequality in (32). Let $d := d(f) \equiv (\int_\Omega 1\wedge |f|^2 d\mu)^{1/2},\ f\in\mathcal{L}^+$ and consider the sets $A=\{|f|\ge 1\}$ and $B=\{|f|<1\}$. Note that $\mu(A)<\infty$ and recall by (7) that $I(f\mathbb{1}_A) = \int_A f d\Pi_\mu - \int_A a(f)d\mu$. From the definition of a and d, see (8) and (11), it follows that $|\int_A a(f)d\mu| \le \mu(A) \le d^2$ and therefore

$$\mathbb{P}\{|I(f\mathbb{1}_A)|>d^2\} \le \mathbb{P}\Big\{\Big|\int_A f d\Pi_\mu\Big| \ne 0\Big\} \le 1-e^{-\mu(A)} \le 1-e^{-d^2}. \tag{37}$$

The second inequality follows from the fact that $\int_A f d\Pi_\mu$ is non–zero only when the Poisson point process Π_μ has at least one point in the set A. Also, $I(f\mathbb{1}_B)$ has (by definition) expectation 0 and variance $\int_B f^2 d\mu \le d^2$. Thus, by the Chebyshev's inequality,

$$\mathbb{P}\{|I(f\mathbb{1}_B)|>d^{2/3}\} \le d^{2/3}. \tag{38}$$

Since $I(f) = I(f\mathbb{1}_A)+I(f\mathbb{1}_B)$, by (37) and (38), in the case $d\le 1$, we get

$$\begin{aligned}\mathbb{P}\{|I(f)|>2d^{2/3}\} &\le \mathbb{P}\{|I(f)|>d^2+d^{2/3}\}\\ &\le \mathbb{P}\{|I(f\mathbb{1}_A)|>d^2\} + \mathbb{P}\{|I(f\mathbb{1}_B)|>d^{2/3}\}\\ &\le 1-e^{-d^2}+d^{2/3}\\ &\le 2d^{2/3}.\end{aligned}$$

Hence $d_{KF}(I(f)) \le 2d^{2/3}$, provided that $d\le 1$. This, since $d_{KF}(I(f))\le 1$ implies the first inequality in (32). $\square$

Proof of Proposition 1.12. The proof is standard. Let $\{f_n\}_{n\in\mathbb{N}} \subset \mathcal{L}^+$ be a Cauchy sequence in d. Then, for all $\varepsilon\in(0,1)$, we have

$$\mu\{|f_m-f_n|>\varepsilon\} \le \frac{1}{\varepsilon^2}\int_\Omega 1\wedge|f_m-f_n|^2 d\mu = \frac{d(f_m,f_n)^2}{\varepsilon^2} \to 0,$$

as $m, n \to \infty$, which shows that $\{f_n\}_{n\in\mathbb{N}}$ is Cauchy in measure. Hence, there exists a sub–sequence $\{n_k\}_{k\in\mathbb{N}}$ and a measurable function f, such that $f_{n_k} \to f$, as $n_k \to \infty$, μ–a.e. Now, by the Fatou's lemma, we obtain

$$d(f_{n_k}, f)^2 = \int_\Omega 1 \wedge |f_{n_k} - f|^2 d\mu \le \liminf_{\ell\to\infty} \int_\Omega 1 \wedge |f_{n_k} - f_{n_\ell}|^2 d\mu = \liminf_{\ell\to\infty} d(f_{n_k}, f_{n_\ell})^2.$$

This inequality implies that $d(f_{n_k}, f) < \infty$, and hence $f \in \mathcal{L}^+$, because $d(f, 0) \le d(f, f_{n_k}) + d(f_{n_k}, 0) < \infty$. Since $\{f_n\}_{n\in\mathbb{N}}$ is Cauchy in the metric d, we also have that $d(f_{n_k}, f) \to 0$, as $n_k \to \infty$, and hence $d(f_n, f) \to 0$, as $n \to \infty$. Thereby proving that the metric d is complete.

Let now $(\Omega, \mathcal{B})$ be Borel. Recall that the measure μ is σ–finite. Then the space $L^2 = L^2(\Omega, \mathcal{B}, \mu)$ $(\subset \mathcal{L}^+)$ equipped with the usual L^2–norm is separable and let $\{f_n\}_{n\in\mathbb{N}}$ be a dense subset of L^2. By (11), for all $f \in \mathcal{L}^+$, $f_n \in L^2$, and $K > 0$, we have

$$\begin{aligned} d(f, f_n)^2 &= \int_\Omega 1 \wedge |f - f_n|^2 d\mu \\ &\le \int_{\{|f|\le K\}} |f - f_n|^2 d\mu + \int_{\{|f|>K\}} d\mu \\ &\le \int_\Omega (f \mathbb{1}_{\{|f|\le K\}} - f_n)^2 d\mu + \mu\{|f| > K\}. \end{aligned}$$

Since $f \in \mathcal{L}^+$, we have that $f \mathbb{1}_{\{|f|\le K\}} \in L^2$ and $\mu\{|f| > K\} \to 0$, as $K \to \infty$. Thus, by picking large enough K and a suitable f_n, one can make $d(f, f_n)$ arbitrarily small, showing that $\{f_n\}_{n\in\mathbb{N}}$ is also dense in the metric space $(\mathcal{L}^+, d)$, thereby proving separability. ☐

Proof of Proposition 1.13. Suppose first that $d(f_n - f) + |c_n - c| \to 0$, as $n \to \infty$. Then, by Slutsky's theorem, it is enough to show that $I(f_n)$ converges in probability to $I(f)$, as $n \to \infty$. By (30), we have that $I(f_n) - I(f) = I(f_n - f) + \gamma(f_n, -f)$. Proposition 3.4 and the assumption $d(f_n - f) \to 0$ imply that $I(f_n - f) \overset{\mathbb{P}}{\to} 0,\ n \to \infty$. It remains to show that $\gamma(f, -f_n) \to 0$, as $n \to \infty$. By the triangle inequality for d, we have $|d(f_n) - d(f)| \le d(f_n - f) \to 0$, as $n \to \infty$, and in particular $d(f_n), n \in \mathbb{N}$ is bounded. Thus, by Lemma 3.2 applied to f and $g := -f_n$, we obtain $\gamma(f, -f_n) \to 0$, as $n \to \infty$. This completes proof of the 'if' part.

To prove the 'only if' part, suppose that $I(f_n) + c_n \overset{\mathbb{P}}{\to} \xi,\ n \to \infty$, set $\xi_{m,n} := I(f_m) - I(f_n) + c_m - c_n$, and let $\xi'_{m,n}$ be and independent copy of $\xi_{m,n}$. Then, by using (9) we obtain that

$$\xi_{m,n} - \xi'_{m,n} \overset{d}{=} I(f_m - f_n) - I(f_m - f_n)',$$

where $I(f_m - f_n)'$ is an independent copy of $I(f_m - f_n)$. Now, by the second bound in (32) of Proposition 3.4 applied to $f := f_m - f_n$, we obtain

$$1 - e^{-cd(f_m - f_n)^2} \le 2d_{KF}(I(f_m - f_n) - I(f_m - f_n)') \equiv 2d_{KF}(\xi_{m,n} - \xi'_{m,n}) \le 4d_{KF}(\xi_{m,n}).$$

The right–hand side of the last inequality vanishes, as $m, n \to \infty$, since the sequence $\{I(f_n) + c_n,\ n \in \mathbb{N}\}$ converges in probability and therefore it is Cauchy in the Ky Fan metric. This implies that $d(f_m - f_n) \to 0,\ m, n \to \infty$, and since $(\mathcal{L}^+, d)$ is complete (Proposition 1.12), there is an $f \in \mathcal{L}^+$, such that $d(f_n - f) \to 0,\ n \to \infty$. Therefore, by the already established 'if' part, it follows that $I(f_n) \overset{\mathbb{P}}{\to} I(f),\ n \to \infty$. This, and the fact that $I(f_n) + c_n \overset{\mathbb{P}}{\to} \xi,\ n \to \infty$ imply (by Slutsky) that the sequence c_n converges to a constant c and $\xi = I(f) + c$. This completes the proof. ☐

Proof of Theorem 1.11. Let T_0 be the at most countable subset of T appearing in Condition S. Consider the space $\mathbb{R}^{T_0}$, equipped with the product σ–algebra $\mathcal{B}$. Following [20] (see also [28]), let μ be the Lévy measure of $\{X(t),\ t \in T_0\}$ on $\mathbb{R}^{T_0}$. For $t \in T_0$, we define the coordinate mappings $f_t : \mathbb{R}^{T_0} \to \mathbb{R}$ by $f_t(\varphi) = \varphi(t)$, where $\varphi : T_0 \to \mathbb{R}$, $\varphi \in \mathbb{R}^{T_0}$. Then, $\{f_t, t \in T_0\}$ is a spectral representation of $\{X(t), t \in T_0\}$ by the properties of the Lévy measure.

For $t \notin T$, observe that by Condition S, there exists a sequence $\{t_n\} \subset T_0$, such that $X(t_n)$ converges in probability to $X(t)$, as $n \to \infty$. With other words, $I(f_n) + c_n$ converges in probability to $I(f) + c$, for some c_n and c. Thus, by Proposition 1.13, the sequence of functions f_{t_n} has a limit in $(\mathcal{L}^+, d)$, as $n \to \infty$. We take this limit to be the spectral function f_t.

Notice that the so–defined spectral representation is minimal. Indeed, the σ–algebra $\sigma\{f_t,\ t \in T\}$ coincides with the product σ–algebra $\mathcal{B}$ on $\mathbb{R}^{T_0}$. We also have that $\operatorname{supp}\{f_t, t \in T_0\} = \mathbb{R}^{T_0}$ (mod μ) because $\cap_{t \in T_0}\{f_t = 0\} = \{0\}$, a set whose Lévy measure is 0 by convention. To complete the proof observe that the measurable space $(\mathbb{R}^{T_0}, \mathcal{B})$ is Borel by Kuratowski's theorem. $\square$

Proof of Theorem 1.14. We are going to apply Lemma 3.1. Define the measurable mappings $F_i : (\Omega_i, \mathcal{B}_i) \to (\mathbb{R}^T, \mathcal{B})$ by

$$F_i(\omega) = (f_t^{(i)}(\omega))_{t \in T}, \quad \omega \in \Omega_i, \quad i = 1, 2.$$

Minimality implies that the first condition of Lemma 3.1 is satisfied. We prove that $\mu_1 \circ F_1^{-1} = \mu_2 \circ F_2^{-1}$. Let $t_1, \dots, t_n \in T$ and observe that in view of (9) we have

$$\begin{aligned}
\mathbb{E} e^{i \sum_{j=1}^n \theta_j X(t_j)} &= \mathbb{E} \exp\Big\{ i \sum_{j=1}^n \theta_j (I(f_{t_j}^{(i)}) + c_j^{(i)}) \Big\} \\
&= \exp\Big\{ i \sum_{j=1}^n c_j^{(i)} \theta_j + \int_{\mathbb{R}^n} \Big(e^{i \sum_{j=1}^n \theta_j x_j} - i \sum_{j=1}^n \theta_j a(x_j) - 1 \Big) (\mu_i \circ G_i^{-1})(dx) \Big\},
\end{aligned}$$

where $G_i = (f_{t_j}^{(i)})_{j=1}^n : \Omega_i \to \mathbb{R}^n$ and $c_1^{(i)}, \dots, c_n^{(i)} \in \mathbb{R}$ are constants, $i = 1, 2$. The last relation and the uniqueness of the Lévy measure of the i.d. random vector $(X(t_j))_{j=1}^n$ shows that $(\mu_1 \circ G_1)^{-1}(A) = (\mu_2 \circ G_2)^{-1}(A)$ for all Borel sets $A \subset \mathbb{R}^n \backslash \{0\}$. We need to show that $(\mu_1 \circ G_1)^{-1}(\{0\}) = (\mu_2 \circ G_2)^{-1}(\{0\})$. As in the proof of Theorem 1.8 we can find a sequence $q_1, q_2, \ldots \in T$ such that $\mu_i(\cap_{j \in \mathbb{N}}\{f_{t_j}^{(i)}\}) = 0$, $i = 1, 2$. Consider measurable sets

$$E_{i,p} = G_i^{-1}(\{0\}) \cap (\cap_{j=1}^{p-1}\{f_{q_j}^{(i)} = 0\}) \cap \{f_{q_p}^{(i)} \neq 0\}.$$

For every p, we have shown that $\mu_1(E_{1,p}) = \mu_2(E_{2,p})$. It follows that

$$\mu_1(G_1^{-1}(\{0\})) = \sum_{p=1}^{\infty} \mu_1(E_{1,p}) = \sum_{p=1}^{\infty} \mu_2(E_{2,p}) = \mu_2(G_2^{-1}(\{0\})).$$

This proves that $(\mu_1 \circ G_1)^{-1}(A) = (\mu_2 \circ G_2)^{-1}(A)$ for all Borel sets $A \subset \mathbb{R}^n$. With other words, the measures $\mu_1 \circ F_1^{-1}$ and $\mu_2 \circ F_2^{-1}$ are equal on the semiring $\mathcal{C}$ consisting of subsets $\{\varphi : T \to \mathbb{R} : (\varphi(t_j))_{j=1}^n \in A\}$, where $A \subset \mathbb{R}^n$ is Borel. This semiring generates the product σ–algebra $\mathcal{B}$. Also, we have a decomposition

$$\mathbb{R}^T = \cup_{n=1}^{\infty} \cup_{k=1}^{\infty} \{\varphi : T \to \mathbb{R} : k^{-1} \le |\varphi(q_n)| \le k\} \quad \text{mod } \mu_1 \circ F_1^{-1} \text{ and } \mu_2 \circ F_2^{-1}.$$

Note that the sets on the right–hand side have finite $\mu_1 \circ F_1^{-1}$ (and $\mu_2 \circ F_2^{-1}$) measure and belong to the semiring $\mathcal{C}$. By the uniqueness of measure extension theorem, the measures $\mu_1 \circ F_1^{-1}$ and $\mu_2 \circ F_2^{-1}$ are equal. Lemma 3.1 completes the proof. □

References

[1] J. Aaronson. *An introduction to infinite ergodic theory*, volume 50 of *Mathematical Surveys and Monographs.* American Mathematical Society, Providence, RI, 1997.

[2] A. A. Balkema, L. de Haan, and R. L. Karandikar. Asymptotic distribution of the maximum of n independent stochastic processes. *J. Appl. Probab.*, 30(1):66–81, 1993.

[3] A. A. Balkema and S. Resnick. Max–infinite divisibility. *J. Appl. Probab.*, 14:309–319, 1977.

[4] L. de Haan. A spectral representation for max–stable processes. *Ann. Probab.*, 12(4):1194–1204, 1984.

[5] L. de Haan and J. Pickands. Stationary min–stable stochastic processes. *Probab. Theory Relat. Fields*, 72:477–492, 1986.

[6] C. Dombry and F. Eyi-Minko. Strong mixing properties of max–infinitely divisible random fields. *Stoch. Processes Appl., to appear*, 2012. Available at http://arxiv.org/abs/1201.4645.

[7] G. Gerritse. Supremum self-decomposable random vectors. *Probab. Theory Relat. Fields*, 72(1):17–33, 1986.

[8] E. Giné, M. G. Hahn, and P. Vatan. Max–infinitely divisible and max–stable sample continuous processes. *Probab. Theory Relat. Fields*, 87(2):139–165, 1990.

[9] P. R. Halmos. *Measure Theory.* Van Nostrand Company, New York, 1950.

[10] C. D. Hardin, Jr. Isometries on subspaces of L^p. *Indiana Univ. Math. J.*, 30:449–465, 1981.

[11] C. D. Hardin, Jr. On the spectral representation of symmetric stable processes. *J. Multivariate Anal.*, 12:385–401, 1982.

[12] Z. Kabluchko. Spectral representations of sum– and max–stable processes. *Extremes*, 12(4):401–424, 2009.

[13] Z. Kabluchko. Stationary systems of Gaussian processes. *Ann. Appl. Probab.*, 20(6):2295–2317, 2010.

[14] Z. Kabluchko. Extremes of independent Gaussian processes. *Extremes*, 14(3):285–310, 2011.

[15] Z. Kabluchko and M. Schlather. Ergodic properties of max-infinitely divisible processes. *Submitted*, 2009. Available at http://arxiv.org/abs/0905.4196.

[16] J. F. C. Kingman. *Poisson processes*, volume 3 of *Oxford Studies in Probability.* The Clarendon Press Oxford University Press, New York, 1993. Oxford Science Publications.

[17] S. Kolodyński and J. Rosiński. Group self–similar stable processes in $\mathbb{R}^d$. *J. Theoret. Probab.*, 16(4): 855–876 (2004), 2003.

[18] G. W. Mackey. Point realizations of transformation groups. *Illinois J. Math.*, 6:327–335, 1962.

[19] D. Maharam. Incompressible transformations. *Fund. Math.*, 56:35–50, 1964.

[20] G. Maruyama. Infinitely divisible processes. *Theory of Probab. Appl.*, 15(1):1–22, 1970.

[21] I. Molchanov. *Theory of random sets.* Probability and its Applications. Springer–Verlag, London, 2005.

[22] M. Penrose. The minimum of a large number of Bessel processes. *J. Lond. Math. Soc., II. Ser.*, 38(3): 566–576, 1988.

[23] M. Penrose. Minima of independent Bessel processes and of distances between Brownian particles. *J. Lond. Math. Soc., II. Ser.*, 43(2):355–366, 1991.

[24] M. Penrose. Semi–min–stable processes. *Ann. Probab.*, 20(3):1450–1463, 1992.

[25] B. S. Rajput and J. Rosiński. Spectral representations of infinitely divisible processes. *Probab. Theory Relat. Fields*, 82(3):451–487, 1989.

[26] J. Rosiński. On the structure of stationary stable processes. *Ann. Probab.*, 23(3):1163–1187, 1995.

[27] J. Rosiński. Decomposition of stationary α–stable random fields. *Ann. Probab.*, 28(4):1797–1813, 2000.

[28] E. Roy. Ergodic properties of Poissonian ID processes. *Ann. Probab.*, 35(2):551–576, 2007.

[29] E. Roy. Maharam extension and stationary stable processes. *Ann. Probab.*, 40(3):1357–1374, 2012.

[30] G. Samorodnitsky. Null flows, positive flows and the structure of stationary symmetric stable processes. *Ann. Probab.*, 33(5):1781–1803, 2005.

[31] K.-I. Sato. *Lévy processes and infinitely divisible distributions.* Cambridge Studies in Advanced Mathematics, 68. Cambridge University Press, 1999.

[32] R. Sikorski. *Boolean algebras.* Third edition. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 25. Springer-Verlag, New York, 1969.

[33] S. Stoev. On the ergodicity and mixing of max–stable processes. *Stoch. Processes Appl.*, 118:1679–1705, 2008.

[34] P. Vatan. Max–infinite divisibility and max–stability in infinite dimensions. Probability in Banach spaces V. Lect. Notes Math. 1153, 400–425., 1985.

[35] Y. Wang, P. Roy, and S. Stoev. Ergodic properties of sum- and max-stable stationary random fields via null and positive group actions. *Submitted*, 2009. Available at http://arxiv.org/abs/0911.0610.

[36] Y. Wang and S. Stoev. On the association of sum– and max–stable processes. *Statistics and Probability Letters*, 80:480–488, 2010.

[37] Y. Wang and S. Stoev. On the structure and representations of max–stable processes. *Adv. Appl. Probab.*, 42(3):855–877, 2010.

Zakhar Kabluchko, Institute of Stochastics, Ulm University, Helmholtzstr. 18, 89069 Ulm, Germany
E-mail address: `zakhar.kabluchko@uni-ulm.de`

Stilian Stoev, Department of Statistics, University of Michigan, Ann Arbor, 439 W. Hall, 1085 S. University Ann Arbor, MI 48109-1107, USA
E-mail address: `sstoev@umich.edu`